

# **Preoperative Patient Information**

©2020 American Society of Plastic Surgeons<sup>®</sup>. Purchasers of the *Informed Consent Resource* are given a limited license to modify documents contained herein and reproduce the modified version for use in the Purchaser's own practice only. All other rights are reserved by the American Society of Plastic Surgeons<sup>®</sup>. Purchasers may not sell or allow any other party to use any version of the *Informed Consent Resource*, any of the documents contained herein, or any modified version of such documents.



## Please fill out the following before surgery.

#### Drug Reactions/Allergies/Latex Sensitivity

Some patients are allergic to medications such as penicillin. Others may get nausea/vomiting from narcotics like codeine, morphine, Demerol<sup>®</sup>, Vicodin<sup>®</sup>, and Percocet<sup>®</sup>.

Please list any known drug allergies, reactions, or sensitivities.

Medicine Name Type of Reaction/Allergy 1. 2. 3. 4. \_\_\_\_\_I do not have any known drug allergies, reactions, or latex sensitivity.

### Prescription Medications

Please list all prescription medications that you are taking.

1. 2. 3.

4.

\_\_I am currently not taking any prescription medications.

### Nonprescription Medications/Dietary Supplements/Vitamins/"Herbs"/Minerals

Many patients take nonprescription medicines. These include aspirin and anti-inflammatories (Advil<sup>®</sup>, Motrin<sup>®</sup>, Aleve<sup>®</sup>). They can also include others like dietary supplements, vitamins, "herbs," or minerals. These may cause bleeding during or after surgery. They may also react with other prescription medications. If you are taking any such items, please list them below. Stop using them for at least 10 days before and after surgery.

1.

2.

3.

\_\_\_\_\_I am currently not taking any nonprescription medications.

#### Smoking, Secondhand Smoke Exposure, Nicotine use

Patients who smoke or use tobacco or nicotine products (patch, gum, or nasal spray) are at risk of surgical problems. This includes skin death and delayed healing. The same is true for people exposed to secondhand smoke or other nicotine products. Smoking can negatively affect recovery from anesthesia. It can cause coughing and increased bleeding. People who are not exposed to nicotine are at a lower risk of these problems. Please mark your current nicotine use below:

\_\_\_\_\_I am a nonsmoker and do not use nicotine products.

\_\_\_\_\_I am a smoker and use tobacco or nicotine products. I know this may cause surgical complications.

\_\_\_\_\_I used to smoke. I stopped about \_\_\_\_\_\_ ago. I know I may still have problems from smoking if not enough time has lapsed.

Date/Time: \_\_\_\_\_ Signature: \_\_\_\_\_

Page 1 of 1

Patient Initials

©2020 American Society of Plastic Surgeons®

This form is for reference purposes only. It is a general guideline and not a statement of standard of care. Rather, this form should be edited and amended to reflect policy requirements of your practice site(s), CMS and Joint Commission requirements, if applicable, and legal requirements of your individual states. The ASPS does not certify that this form, or any modified version of this form, meets the requirements to obtain informed consent for this procedure in the jurisdiction of your practice.